

A Note on TeV Cerenkov Events as Bose-Einstein Gamma Condensations

Amir Levinson

School of Physics and Astronomy, Tel Aviv University, Tel Aviv 69978, Israel

ABSTRACT

The idea that the TeV air showers, thought to be produced by > 10 TeV gamma rays from Mrk 501, can be mimicked by coherent bunches of sub-TeV photons is reexamined, focusing on fundamental considerations. In particular, it is shown that the minimum spot size of the beam of pulsed TeV photons arriving at Earth is on the order of a few kilometers, unless a lens with certain characteristics is placed between the TeV laser and Earth. The viability of the laser production mechanism proposed by Harwit et al. (2000) is also reassessed.

It has been argued recently (e.g., Coppi & Aharonian 1999) that the detection of TeV photons from Mrk 501 at energies well above 10 TeV (Hayashida, et al. 1998; Aharonian et al. 1999) places severe constraints on the diffuse extragalactic IR background, and may be particularly problematic (Protheroe & Meyer, 2000) in view of recent determinations of the IR background by various experiments. Motivated by this consideration, Harwit et al. (1999) proposed that the observed air showers, that are commonly interpreted as due to single > 10 TeV gamma rays, can be produced by coherent bunches of sub-TeV photons that are not absorbed by pair production on the extragalactic IR background. What they envisaged essentially is that some mechanism in the source (Mrk 501) produces pulsed, sub-TeV lasers that, when interacting with the Earth's atmosphere could mimic TeV air shower events. This hypothesis has been tested subsequently by the HEGRA collaboration (Aharonian et al. 2000), who claimed that it can be rejected on the basis of a comparison of the energy dependent penetration depth in Earth's atmosphere of TeV photons from Mrk 501 with the penetration depth of photons from the Crab Nebula. In view of the growing interest in this scenario it is worth reexamining its viability. Below, we discuss the fundamental limitations of such a TeV laser.

What are the basic requirements from the system under consideration?

i) A typical Cerenkov flash produced by a TeV air shower lasts for about several ns. This implies that the width of the TeV pulse produced by the laser, Δt , should not exceed this timescale, and that temporal coherence must be maintained over a time $> \Delta t$. The corresponding light crossing time, $c\Delta t$, is of the order of several meters. In principle, however, the dimension of the system should not be restricted to this scale. In laboratory lasers, for instance, pulse durations as short as the decay time of the lasing substrate (which can be shorter by many orders of magnitude than the light crossing time of the cavity, though typically larger than the beam diameter) can be achieved using e.g., mode locking or Q switching methods (which require modulation of either the pumping rate or the refraction index in the cavity) (e.g., Svelto 1998). Although it is difficult to envisage how this situation can be accomplished under astrophysical conditions, the requirement that the size of the system would not exceed the pulse width does not seem to be fundamental. Moreover, if the laser mechanism involves relativistic motion the pulse can be further compressed, owing to time dilation effects.

ii) The spot size of the TeV laser beam illuminating Earth should be within the angular resolution of current TeV experiments, otherwise the shower image will differ from that expected to be produced by a single TeV photon. (If the spot is resolved, it can give rise to a shower image that may resemble that of a

cosmic ray shower. Such an event is likely to be rejected.) For a typical angular resolution of 0.1° , and shower height of say 10 km, this yields a spot size < 20 meters. As shown below this requirement places a stringent constraint on the system.

Harwit et al. argued that the arriving bunched photons would spread over a distance $\Delta y \simeq h/\Delta p_y \simeq \lambda L/D$, where Δp_y is the uncertainty in transverse momentum, which equals approximately the product of the photon momentum, h/λ , and the angular size of the source subtends at Earth, D/L , with D being the characteristic source dimension and L the distance between Earth and the source (Mrk 501). Taking D to be on the order of a Schwarzschild radius of a supermassive black hole yields Δy in the range between 10^{-1} to 10^{-4} cm for the expected range of black hole masses, which, as they argued, is negligible in Cerenkov detection. It should be noted, however, that Δy is essentially the size of the diffraction wing of a beam having a diameter D at a distance L from the beam waist (see Fig. 1) and not the spot size, which for the above parameters is practically D . Thus, if the dimension of the photon bunch (or Bose Einstein condensate, as termed by those authors) is indeed $D \sim r_g$, then in the absence of an additional optical system (see below), the arriving photon bunch would spread over a similar scale, not $\lambda L/r_g$. We suspect though that the picture envisioned by Harwit et al. is that each condensate is well localized, so that the spatial dimension of each bunch at the source is not D but very small. In that case the uncertainty in transverse momentum of a photon in the bunch is related to the latter scale and is much larger than that quoted above. The question then is, what is the minimum spot size that can be achieved given the distance between Earth and Mrk 501?

Consider some apparatus that produces a pulsed TeV laser beam having a diameter D at the beam waist (see Fig. 1). The diffraction angle of the beam is $\psi = \lambda/D$, where $\lambda = 2 \times 10^{-17}(\epsilon/1\text{TeV})^{-1}$ cm is the wavelength of the laser at its spectral peak, and ϵ is the corresponding energy. At a distance L from the laser the beam spot size, a , is the sum of the waist spot size and the size of the diffraction wing:

$$a = D + \psi L = D + (\lambda/D)L. \quad (1)$$

For a target at a fixed distance L from the laser, the minimum beam spot size a_{min} can be obtained by minimizing a with respect to D ; that is taking $da/dD = 0$. This yields $D = \sqrt{\lambda L}$ and

$$a_{min} = 2\sqrt{\lambda L} \simeq 10^5 (L/100\text{Mpc})(\epsilon/1\text{TeV})^{-1} \text{ cm}. \quad (2)$$

Taking L to be the distance from Earth to Mrk 501 ($L = 130$ Mpc assuming $h_o = 0.65$), we conclude that the arriving pulse of sub TeV photons would spread over a distance of at least several kilometers.

The spot size can in principle be reduced if a lens is placed between the source and Earth. In the optimal case the radius of the lens should be comparable to the beam radius, and its focal plan should




Fig. 1.— Schematic illustration of the propagation of a coherent TeV beam having a diameter D at its waist. The diffraction angle of the beam is $\psi = \lambda/D$, where λ is the corresponding wavelength, and the beam spot size a distance L away is $a = D + \psi L$.

intersect Earth. Under such conditions the size of the spot illuminated by the TeV laser is diffraction limited (assuming complete coherence). Denoting by D_l the lens diameter and by L_l its distance from Earth, and requiring a spot size smaller than say 10 meters, yields a minimum lens diameter of

$$D_{lmin} = 10^7 (L_l/100\text{Mpc})(\epsilon/1\text{TeV})^{-1} \text{ cm.} \quad (3)$$

This is larger than the gravitational radius of a stellar mass object. As an illustrative example consider lensing of the TeV beam by a point mass located on the axis of the beam a distance L_l from Earth. In that case only rays for which the impact parameter lies in the range between b_+ and b_- , where $b_{\pm} \simeq 2\sqrt{r_g L_l} \pm d$, d being the diameter of the beam spot on Earth, and r_g is the gravitational radius of the lens, will be deflected by the required angle. For typical parameters, this constitutes only a tiny fraction of the beam photons and, therefore, a point mass cannot provide the required lens. What seems to be needed is some extended object with a density profile such that the refraction index of the lens would be independent of the lens radius. Perhaps high velocity clouds?

We conclude by briefly commenting on the TeV laser production mechanism discussed by Harwit et al. These authors suggested that inverse Compton scattering of OH or H₂O megamaser photons by a relativistic beam of nonthermal (in the comoving frame) electrons, may provide a mean for producing a coherent TeV pulse. We stress that spontaneous scattering is a process that tends to destroy coherence; a single seed photon has a probability of being scattered into many different directions with different phases. (It is worth noting that the occupation number of scattered photons will be reduced not only by the ratio of bandwidths, as pointed out by Harwit et al., but also by the ratio of angular sizes of the beam of scattered photons and the megamaser beam, which can in principle be large.) The fact that the occupation number may largely exceed unity does not imply that the emission is coherent, unless the phases of emitted photons are somehow correlated. In order to mimic TeV air showers, it is required that several photons will be scattered into a solid angle, henceforth denoted by $\Delta\Omega$, not larger than that subtends at the source by the maximum spot size allowed in the Earth's atmosphere, and will have a time spread of no more than several ns. Since the scattering process is unsynchronized, it implies that the characteristic dimension of the emitting volume should not exceed $\Gamma^2 c \Delta t$, where $\Delta t \sim 10$ ns is again the time scale of a typical Cerenkov shower, and Γ is the bulk Lorentz factor of the electron beam. Using the number density of maser photons adopted by Harwit et al., $n_{ph} \sim 10^{10} \text{ cm}^{-3}$, yields a scattering rate per electron of $\nu_{sc} = \sigma_T c n_{ph} \sim 2 \times 10^{-4} \text{ s}^{-1}$. The total scattering rate by a beam section of volume $\Delta V \sim (\Gamma^2 c \Delta t)^3$ containing relativistic electrons having density n_e is then $\nu_{sc} \Delta V n_e \sim 2 \times 10^{16} (\Delta t/10 \text{ ns})^3 (\Gamma/100)^6 (n_e/10^2 \text{ cm}^{-3}) \text{ s}^{-1}$. Now, only a fraction $\Delta\Omega/\Delta\Omega_b$ will be scattered into a solid angle $\Delta\Omega$, where $\Delta\Omega_b$ denotes the solid angle of the TeV photon beam, which in the case of scattering by an electron beam having bulk Lorentz factor Γ is on the order of $\Delta\Omega_b \sim \pi\Gamma^{-2}$. For a maximum spot size d we thus obtain: $\Delta\Omega/\Delta\Omega_b \simeq 10^{-43} (\Gamma/100)^2 (d/10^3 \text{ cm})^2 (L/100 \text{ Mpc})^{-2}$. Consequently, the rate at which the maser photons in the volume ΔV are scattered into a given direction within a solid angle $\Delta\Omega$ is approximately 10^{-27} s^{-1} , about 35 orders of magnitudes smaller than what required.

I thank M. Segev, D. Eichler, D. Maoz, and A. Gal Yam for enlightening discussions. Support from the Israel Science Foundation is greatly acknowledged.

REFERENCES

- Aharonian, F.A. et al., 1999, A&A, 349, 11
 Aharonian, F. et al., 2000 (astro-ph/0006092)

- Coppi, P.S. & Aharonian, F.A. 1999, *Aph*, 11, 35
- Harwit, M., Protheroe, R.J. and Biermann, P.L. 1999, *ApJ*, 524, L91
- Hayashida, N., et al. 1998, *ApJ*, 504, L71
- Protheroe, R.J. and Meyer, H. 2000, *Phys. Lett.*, submitted (astro-ph/0005349)
- Svelto, O., 1998, *Principles of Lasers*, (Plenum Press: NY)